

For Research Use Only

# G3BP2 Monoclonal antibody, PBS Only

Catalog Number: 68580-1-PBS

Featured Product



## Basic Information

<b>Catalog Number:</b> 68580-1-PBS	<b>GenBank Accession Number:</b> BC011731	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 100ug, Concentration: 1mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 9908	<b>CloneNo.:</b> 2E5G3
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> Q9UN86	
<b>Isotype:</b> IgG1	<b>Full Name:</b> GTPase activating protein (SH3 domain) binding protein 2	
<b>Immunogen Catalog Number:</b> AG9222	<b>Calculated MW:</b> 482aa,54 kDa; 449aa,51 kDa	
	<b>Observed MW:</b> 54 kDa	

## Applications

**Tested Applications:**  
WB, IF/ICC, Indirect ELISA

**Species Specificity:**  
human, rat

## Background Information

Stress granules (SGs) are cytoplasmic mRNA-protein condensates formed in response to cellular stressors, such as oxidative stress, ultraviolet radiation, and viral infection (1). The Ras-GTPase-activating protein-binding proteins (G3BPs), consisting of G3BP1 and G3BP2, are key nucleating factors essential for SG formation. They function to protect RNAs from harmful conditions. G3BP2 is mainly distributed in the cytoplasm and participates in the formation of stress granules, cell differentiation, proliferation, and signal transduction. Accumulating evidence has demonstrated that aberrant expression of G3BP2 contributes to cancer initiation and progression, such as high expression of G3BP2 increasing cell stemness, metastasis and chemoresistance in breast cancer.

## Storage

**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed sodium arsenite treated HeLa cells using G3BP2 antibody (68580-1-Ig, Clone: 2E5G3) at dilution of 1:800 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 68580-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 68580-1-Ig (G3BP2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 68580-1-PBS in a different storage buffer formulation.



WB result of G3BP2 antibody (68580-1-Ig; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-G3BP2 transfected HepG2 cells. This data was developed using the same antibody clone with 68580-1-PBS in a different storage buffer formulation.