

For Research Use Only

# TOX2 Monoclonal antibody, PBS Only



Catalog Number: 68590-1-PBS

## Basic Information

<b>Catalog Number:</b> 68590-1-PBS	<b>GenBank Accession Number:</b> NM_001098797	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 100ug , Concentration: 1 mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 84969	<b>CloneNo.:</b> 2G12A7
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> Q96NM4	
<b>Isotype:</b> IgG1	<b>Full Name:</b> TOX high mobility group box family member 2	
<b>Immunogen Catalog Number:</b> AG31563	<b>Calculated MW:</b> 52KD	
	<b>Observed MW:</b> 70 kDa	

## Applications

**Tested Applications:**  
WB, Indirect ELISA

**Species Specificity:**  
Human, mouse, rat, rabbit

## Background Information

TOX2 is a transcription factor belonging to the TOX (thymocyte selection-associated HMG box) family that shares a highly conserved high mobility group DNA-binding domain with the other TOX members. A previous study showed that TOX2 is preferentially expressed in human NK cells among various leukocyte populations and is required for in vitro and in vivo human NK cell differentiation from UCB-derived CD341 hematopoietic stem cells (HSCs) (PMID: 25352127).

## Storage

**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68590-1-Ig (TOX2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 68590-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 68590-1-Ig (TOX2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 68590-1-PBS in a different storage buffer formulation.