For Research Use Only

STAT2 Recombinant antibody, PBS Only

Catalog Number:80061-5-PBS

proteintech®
Antibodies | ELISA kits | Proteins
Uni-rAb
www.ptglab.com

Basic Information

Catalog Number:
80061-5-PBS

Size:
100ug , Concentration: 1 mg/ml by Nanodrop;

Source:
Rabbit

Isotype:
IgG

Immunogen Catalog Number:
AG10168

GenBank Accession Number:
BC051284

GeneID (NCBI):
6773

UNIPROT ID:
P52630

Full Name:
signal transducer and activator of transcription 2, 113kDa

Calculated MW:
851 aa, 98 kDa

Observed MW:
110 kDa

Purification Method:
Protein A purification

CloneNo.:
242685A1

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human

Background Information

STAT2, also named as p113, belongs to the transcription factor STAT family. It is a signal transducer and activator of transcription that mediates signaling by type I IFNs (IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jak kinases (TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize and associate with ISGF3G/IRF-9 to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element (ISRE) to activate the transcription of IF stimulated genes, which drive the cell in an antiviral state. It also interacts with CRSP2, CRSP6, Simian virus 5 protein V, rabies virus phosphoprotein, IFNAR1 and IFNAR2. Its interaction with dengue virus NS5 inhibits the phosphorylation of STAT2, and, when all viral proteins are present (polyprotein), STAT2 is targeted for degradation. The calculated molecular weight of STAT2 is 98 kDa, but phosphorylated STAT2 is about 100-113 kDa.

Storage

Storage:
Store at -80°C.
Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 80061-5-RR (STAT2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80061-5-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 80061-5-RR against Human STAT2 were performed. The affinity constant is below 1 pM.