

For Research Use Only

# P53 Recombinant antibody

Catalog Number: 80077-1-RR **Featured Product**



## Basic Information

<b>Catalog Number:</b> 80077-1-RR	<b>GenBank Accession Number:</b> BC003596	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ul , Concentration: 500 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 7157	<b>CloneNo.:</b> 2H13
<b>Source:</b> Rabbit	<b>Full Name:</b> tumor protein p53	<b>Recommended Dilutions:</b> WB 1:5000-1:50000 IF 1:50-1:500
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 44 kDa	
	<b>Observed MW:</b> 53 kDa	

## Applications

**Tested Applications:**  
IF, WB, ELISA

**Species Specificity:**  
Human, mouse, rat, zebrafish

**Positive Controls:**

**WB :** MCF-7 cells, SW480 cells, HCT 116 cells, HT-29 cells, A431 cells, ROS1728 cells, HSC-T6 cells, ATDC-5 cells, Neuro-2a cells, zebrafish tissue, COLO 320 cells, MDA-MB-231 cells, HT-29 cells

**IF :** HepG2 cells,

## Background Information

TP53, also known as P53 and NY-CO-13, belongs to the p53 family and has 9 isoforms. In SDS-Page, the observed molecular weight is about 53 kDa. TP53 acts as a tumor suppressor in many tumor types, including growth arrest or apoptosis depending on the physiological circumstances and cell types. It is involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. TP53 Localizes in the nucleus in most cells but found in the cytoplasm in some cells. (PMID: 26166714; PMID: 25225161)

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.


**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

**Aliquoting is unnecessary for -20°C storage**


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 80077-1-RR (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using P53 antibody (80077-1-RR, Clone: 2H13) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




zebrafish tissue were subjected to SDS PAGE followed by western blot with 80077-1-RR (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 80077-1-RR (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 80077-1-RR (P53 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



WB result of P53 antibody (80077-1-RR; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-P53 transfected A431 cells.