For Research Use Only

PEX14 (human specific) Recombinant antibody

proteintech www.ptglab.com

Catalog Number:80196-1-RR

Basic Information

Catalog Number: GenBank Accession Number:

80196-1-RR BC006327 GeneID (NCBI):

100ul , Concentration: 700 $\mu g/ml$ by 6K5 Nanodrop: **UNIPROT ID:**

075381 Rabbit Full Name: Isotype: peroxisomal biogenesis factor 14

IgG

Calculated MW: Immunogen Catalog Number: 41 kDa

AG0932 Observed MW:

57 kDa

Applications

Tested Applications: FC, IF, IHC, WB, ELISA

Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Protein A purification

CloneNo.:

Positive Controls:

IF: HeLa cells,

Recommended Dilutions:

WB 1:2000-1:14000 IHC 1:200-1:800 IF 1:50-1:500

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat

IHC: human liver cancer tissue, human kidney tissue

Background Information

PEX14 (peroxisomal biogenesis factor 14) is a peroxisomal membrane protein that is essential for protein docking onto the peroxisomes. It is a central component of the peroxisimal matrix protein import machinery and interacts with PEX5 and PEX19. PEX14 is ubiquitously expressed and defects in PEX14 are the cause of peroxisome biogenesis disorder complementation group K (PBD-CGK). This antibody can be used to detect endogenous PEX14 with an apparent molecular weight of 57 kDa (PMID: 16449325; 9653144) and recognize peroxisomal structures in human,

This antibody is specific to human PEX14. It can't detect mouse or rat PEX14.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 80196-1-RR (PEX14 (human specific) antibody) at dilution of 1:7000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using PEX14 antibody (80196-1-RR, Clone: 6K5) at dilution of 1:150 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 80196-1-RR (PEX14 (human specific) antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 80196-1-RR (PEX14 (human specific) antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human PEX14 (human specific) (80196-1-RR, Clone:6K5) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).