

For Research Use Only

Phospho-STAT3 (Ser727) Recombinant antibody

Catalog Number: 80199-2-RR



Basic Information

Catalog Number: 80199-2-RR	GenBank Accession Number: BC000627	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 6774	CloneNo.: 240778A12
Source: Rabbit	UNIPROT ID: P40763	Recommended Dilutions: WB 1:1000-1:4000
Isotype: IgG	Full Name: signal transducer and activator of transcription 3 (acute-phase response factor)	
	Calculated MW: 770 aa, 88 kDa	
	Observed MW: 100 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : HeLa cells,
Species Specificity: human	

Background Information

Signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3, synonyms: APRF, FLJ20882, MGC16063) is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT3 is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. STAT3 mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1×10^6 Calyculin A treated HEK-293T cells were intracellularly stained with 0.25 ug Stat3 Recombinant Antibody (80199-2-RR, Clone:240778A12) and PE-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.



Non-treated HeLa cells, phosphatase inhibitor treated HeLa cells and λ phosphatase treated HeLa cells were subjected to SDS PAGE followed by western blot with 80199-2-RR (Phospho-STAT3 (Ser727) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.