## For Research Use Only

## PTEN Recombinant antibody

Catalog Number:80718-7-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

80718-7-RR BC005821 GeneID (NCBI):

100ul, Concentration: 1000 ug/ml by 5728

Nanodrop: **UNIPROT ID:** P60484

Rabbit Full Name:

Isotype: phosphatase and tensin homolog

IgG Calculated MW: Immunogen Catalog Number: 47 kDa

AG17274

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra), ELISA

Species Specificity:

**Purification Method:** Protein A purfication

CloneNo.:

240134F7

Recommended Dilutions: IF/ICC 1:150-1:600

Positive Controls:

IF/ICC : HeLa cells,

## **Background Information**

PTEN is one of the most critical tumor suppressors, which functions at different subcellular locations, including the plasma membrane and nucleus. The PTEN protein is located at different subcellular regions-PTEN at the plasma membrane suppresses PI3-kinase signaling in cell growth, whereas PTEN in the nucleus maintains genome integrity. At the plasma membrane, PTEN counteracts PI3 kinase signaling by dephosphorylating the potent second messenger PIP3 to PIP2. The loss of PTEN in cancer cells results in over-activation of AKT and mTOR signaling, leading to excessive stimulation of cell growth and inhibition of cell death. In the nucleus, PTEN functions in DNA repair, genome stability, and cell cycle control through associations with Rad51 and p53. PTEN stability is primarily regulated by phosphorylation of C-terminal tail domains (Thr366, Ser370, Ser380, Thr382, Thr383, and Ser385). The phosphorylation leads to a "closed" state of PTEN and maintains PTEN stability. Dephosphorylation of the Cterminal tail opens the PTEN phosphatase domain, thereby increasing PTEN activity. PTEN protein is of the apparent molecular mass expected for PTEN (55 kDa) and PTENa (70 kDa). (PMID: 33083717, PMID: 20622047, PMID: 24768297)

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PTEN antibody (80718-7-RR, Clone: 240134F7) at dilution of 1:300 and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-Phalloidin (red).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug PTEN Recombinant antibody (80718-7-RR, Clone:240134F7) and APC-Conjugated Goat Anti-Rabbit 1gG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLL) kinetic assays of 80718-7-RR against Human PTEN were performed. The affinity constant is below 1 pM.