For Research Use Only

## ubiquitin Recombinant antibody, PBS Only

Catalog Number:80992-1-PBS



**Basic Information** 

Catalog Number: 80992-1-PBS

GenBank Accession Number:

BC000379

GeneID (NCBI):

100ug , Concentration: 1mg/ml by 7314

Nanodrop; UNIPROT ID:
Source: POCG47
Rabbit Full Name:
Isotype: ubiquitin B

IgG

Immunogen Catalog Number:

AG0260

Purification Method: Protein A purification

CloneNo.:

6H6

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity:

human, mouse, rat, hamster, yeast, dog, spinach


## **Background Information**

Ubiquitin B (UBB) is a member of ubiquitin family, one of the most conserved proteins known. Ubiquitin B is required for ATP-dependent, non-lysosomal intracellular protein degradation of abnormal proteins and normal proteins with a rapid turnover. Ubiquitin B is covalently bound to proteins to be degraded, and presumably labels these proteins for degradation. Ubiquitin also binds to histone H2A in actively transcribed regions but does not cause histone H2A degradation, suggesting that ubiquitin is also involved in regulation of gene expression. When polyubiquitin is free (unanchored-polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling. This gene consists of three direct repeats of the ubiquitin coding sequence with no spacer sequence. Consequently, the protein is expressed as a polyubiquitin precursor with a final amino acid after the last repeat. Aberrant form of this protein has been noticed in patients with Alzheimer's and Down syndrome. Interestingly ubiquitin also becomes covalently bonded to many types of pathological inclusions which appear to be resistant to normal degradation.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 80992-1-RR (ubiquitin antibody) at dilution of 1:60000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80992-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 80992-1-RR (ubiquitin antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80992-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 80992-1-RR (ubiquitin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80992-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using ubiquitin antibody (80992-1-RR, Clone: 6H6) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red). This data was developed using the same antibody clone with 80992-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLL) kinetic assays of 80992-1-RR against Human ubiquitin were performed. The affinity constant is 1.57 nM.