

For Research Use Only

# HA Tag Recombinant antibody, PBS Only

Catalog Number: 81290-1-PBS



## Basic Information

Catalog Number:

81290-1-PBS

Size:

100ug , Concentration: 1mg/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

GeneID (NCBI):

6

Full Name:

HA Tag

Purification Method:

Protein A purification

CloneNo.:

6I21

## Applications

Tested Applications:

WB, IF/ICC, IP, Indirect ELISA

Species Specificity:

recombinant protein

## Background Information

Protein tags are a protein or peptide sequences located either on the C- or N- terminal of the target protein, which can facilitate solubility, detection, purification, localization, and expression of the target protein. The HA tag corresponds to amino acid residues YPYDVPDYA of a surface glycoprotein -human influenza virus hemagglutinin (HA). The HA tag is commonly used for a variety of research applications including chromatin immunoprecipitation, ELISA, flow cytometry, western blotting, immunocytochemistry/immunofluorescence among of others.

## Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




HA-tag recombinant protein, non-transfected HEK-293 cells, transfected HEK-293 cells were subjected to SDS PAGE followed by western blot with 81290-1-RR (HA Tag antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 81290-1-PBS in a different storage buffer formulation.




IP result of anti-HA Tag (IP:81290-1-RR, 4ug; Detection:81290-1-RR 1:10000) with Transfected HEK-293T cells lysate 400 ug. This data was developed using the same antibody clone with 81290-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed Transfected HEK-293 cells using HA Tag antibody (81290-1-RR, Clone: 6l21) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 81290-1-PBS in a different storage buffer formulation.



1x10<sup>6</sup> Transfected HEK-293 cells were intracellularly stained with 0.25 ug HA Tag Recombinant antibody (81290-1-RR, Clone:6l21) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), and 0.25 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 81290-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed Transfected HEK-293 cells using HA Tag antibody (81290-1-RR, Clone: 6l21) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 81290-1-PBS in a different storage buffer formulation.