

For Research Use Only

VDAC1/Porin Recombinant antibody



Catalog Number: 81538-1-RR **1 Publications**

Basic Information

Catalog Number: 81538-1-RR	GenBank Accession Number: NM_003374	Purification Method: Protein A purification
Size: 100ul , Concentration: 500 µg/ml by Nanodrop;	GeneID (NCBI): 7416	CloneNo.: 2F4
Source: Rabbit	UNIPROT ID: P21796	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:500-1:2000 IF 1:50-1:500
Isotype: IgG	Full Name: voltage-dependent anion channel 1	
	Calculated MW: 31 kDa	
	Observed MW: 31 kDa	

Applications

Tested Applications: WB, IF, IHC, ELISA	Positive Controls: WB : MDA-MB-231 cells, HeLa cells, LNCaP cells, Jurkat cells, K-562 cells, NIH/3T3 cells, HSC-T6 cells
Cited Applications: WB	IHC : mouse liver tissue,
Species Specificity: Human, mouse, rat	IF : human liver tissue,
Cited Species: mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

VDAC1, also named as VDAC, porin 31HM, porin 31HL and plasmalemmal porin, belongs to the eukaryotic mitochondrial porin family. It adopts an open conformation at low or zero membrane potential and a closed conformation at potentials above 30-40 mV, to form a channel through the mitochondrial outer membrane and also the plasma membrane. Unlike other membrane transport proteins, porins are large enough to allow passive diffusion. Studies have shown that VDAC1 is subject to both phosphorylation and acetylation (PMID: 23233904). The apparent molecular weight of VDAC1 is 30-37 kDa (PMID: 14573604; 23754752; 25681439). Hypoxic conditions were found to trigger cleavage of the VDAC1 C-terminal to yield a 26-kDa truncated but active form (PMID: 22389449; 23233904). This antibody is specific to VDAC1.

Notable Publications

Author	Pubmed ID	Journal	Application
Jingjing Liu	36587874	J Nutr Biochem	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81538-1-RR (VDAC1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 81538-1-RR (VDAC1/Porin antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human liver tissue using VDAC1/Porin antibody (81538-1-RR, Clone: 2F4) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).