For Research Use Only

Phospho-4EBP1 (Thr37) Recombinant antibody

Catalog Number:81812-4-RR




Basic Information	Catalog Number: 81812-4-RR	GenBank Accession Number: BC004459	Purification Method: Protein A purification
	100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GenelD (NCBI): 1978	CloneNo.: 240535G3
		UNIPROT ID: Q13541	Recommended Dilutions: WB 1:1000-1:4000
		Full Name: eukaryotic translation initiation factor 4E binding protein 1 Calculated MW: 118 aa, 12 kDa	
		Applications	Tested Applications: WB, ELISA
Species Specificity: human	WB : HEK-293T cells, λ phosphatase treated HEK-293T cells		
Background Information	4EBP1 (4E-BP1) is a translational regulator and functions by sequestering eukaryotic translation initiation factor 4E (eIF4E) from the translation initiation machinery. 4E-BP1 contains at least six phosphorylation sites (Thr37, Thr46, Ser65, Thr70, Ser83, and Ser112), four of which (Thr37, Thr46, Ser65, and Thr70) are known to be regulated by mTOR signaling. Phosphorylation of Thr37 and Thr46 is thought to prime 4E-BP1 for sequential phosphorylation of Ser65 and Thr70, which results in the dissociation of 4E-BP1 from eIF4E. (PMID: 12747827)		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer. PBS with 0.02% sodium azide and 50% glycerol pH 7.3.		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	0, 1	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated HEK-293T cells, phosphatase inhibitor treated HEK-293T cells and λ phosphatase treated HEK-293T cells were subjected to SDS PAGE followed by western blot with 81812-4-RR (Phospho-4EBP1 (Thr37) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin (66031-1-lg) antibody as a loading control.