

For Research Use Only

Phospho-PERK/EIF2AK3 (Thr982) Recombinant antibody, PBS Only

Catalog Number: 82534-1-PBS



Basic Information

Catalog Number:

82534-1-PBS

Size:

100ug, Concentration: 1mg/ml by
Nanodrop;

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC126354

GeneID (NCBI):

9451

UNIPROT ID:

Q9NZJ5

Full Name:

eukaryotic translation initiation
factor 2-alpha kinase 3

Calculated MW:

1116 aa, 125 kDa

Observed MW:

180 kDa

Purification Method:

Protein A purification

CloneNo.:

4E16

Applications

Tested Applications:

WB, FC (Intra), Indirect ELISA

Species Specificity:

human, mouse

Background Information

EIF2AK3 encodes the protein kinase RNA-like ER kinase (PERK), a key regulator of the unfolded protein response (UPR) in response to ER stress. Under ER stress conditions, activation of PERK is triggered by the dissociation of glucose-regulated protein (GRP) 78 (also known as BiP) from its luminal domain, followed by oligomerization and autophosphorylation. Phosphorylated PERK subsequently phosphorylates eukaryotic translation initiation factor 2 alpha (eif2a), to attenuate global protein translation and reduce incoming ER protein load via upregulated ER chaperone expression. (PMID: 35922637, PMID: 32029570)

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Non-treated NIH/3T3 cells and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 82534-1-RR (Phospho-PERK/EIF2AK3 (Thr982) antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82534-1-PBS in a different storage buffer formulation.



Non-treated HEK-293 cells and Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 82534-1-RR (Phospho-PERK/EIF2AK3 (Thr982) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with beta actin antibody (66009-1-Ig) as loading control. This data was developed using the same antibody clone with 82534-1-PBS in a different storage buffer formulation.



1X10⁶ HEK-293 cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.13 ug Phospho-PERK/EIF2AK3 (Thr982) Recombinant antibody (82534-1-RR, Clone:4E16) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.13 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH. This data was developed

