## For Research Use Only

## YY1 Recombinant antibody

Catalog Number:82712-5-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

82712-5-RR BC037308 GeneID (NCBI):

100ul, Concentration: 700 ug/ml by 7528 2K12

Nanodrop: UNIPROT ID: P25490 Rabbit Full Name:

Isotype: YY1 transcription factor

IgG Calculated MW: Immunogen Catalog Number: 414 aa, 45 kDa

AG17792

**Applications** 

**Tested Applications:** IF/ICC, FC (Intra), ELISA

Species Specificity:

**Purification Method:** 

Protein A purification

CloneNo.:

**Positive Controls:** 

IF/ICC: HepG2 cells,

Recommended Dilutions: IF/ICC 1:350-1:1400

## **Background Information**

YY1, also named as DELTA, INO 80S and NF-E1, contains four C2H2-type zinc fingers and belongs to the YY transcription factor family. YY1 is a multifunctional transcription factor that exhibits positive and negative control on a large number of cellular and viral genes by binding to sites overlapping the transcription start site. YY1 may direct histone deacetylases and histone acetyltransferases to a promoter in order to activate or repress the promoter, thus implicating histone modification in the YY1. The open reading frame of the human YY1 cDNA encodes a protein of 414 amino acids with a predicted molecular weight of 44 kDa. However, YY1 migrates on SDS gels as a 65-68 kDa protein, probably due to the structure of the protein. It is a ubiquitously expressed transcription factor with fundamental roles in embryogenesis, differentiation, replication and proliferation.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Aliquoting is unnecessary for -20°C storage

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

\*\*\* 20ul sizes contain 0.1% BSA


## Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using YY1 antibody (82712-5-RR, Clone: 2K12) at dilution of 1:400 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using YY1 antibody (82712-5-RR, Clone: 2K12 ) at dilution of 1:700 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug YY1 Recombinant antibody (82712-5-RR, Clone:2K12) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (5A00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLL) kinetic assays of 82712-5-RR against Human YY1 were performed. The affinity constant is 0.412 pM.