For Research Use Only

SIRT1 Recombinant antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

2F11

Catalog Number:82790-4-PBS

Basic Information

Catalog Number:

82790-4-PBS

100ug, Concentration: 1 mg/ml by

Nanodrop:

Rabbit Isotype: IgG

Immunogen Catalog Number:

AG3808

GenBank Accession Number:

BC012499 GeneID (NCBI):

23411 **UNIPROT ID:**

Q96EB6 Full Name:

sirtuin (silent mating type

information regulation 2 homolog) 1

(S. cerevisiae) Calculated MW: 747 aa, 82 kDa

Observed MW: 110-130 kDa

Applications

Tested Applications:

WB. IF. Indirect ELISA Species Specificity:

Human

Background Information


SIRT1, also named as SIR2L1, contains a deacetylase sirtuin-type domain and belongs to the sirtuin family. The posttranslation modified SIRT1 is a 110-130 kDa protein, which contains one deacetylase sirtuin-type domain. The 75-80 kDa SirT1 fragment was detected to lack the carboxy-terminus (PMID:21305533). SirT1 exists a 57-61 kDa isoform. SIRT1 may be found in nucleolus, nuclear euchromatin, heterochromatin, and inner membrane. It can shuttles $between the nucleus and cytoplasm. SIRT1 \, regulates \, processes \, such as \, apoptosis \, and \, muscle \, differentiation \, by \, differentiation \, by \, differentiation \, diff$ deacetylating key proteins. SIRT1 in particular initiates several signaling events relevant to cardioprotection, including activation of endothelial nitric oxide synthase, INS receptor signaling, and autophagy. In addition, SIRT1 activation elicits resistance to oxidative stress via the regulation of transcription factors and co-activators such as FOXO, Hif-2a, and NF-kB. SIRT1 regulates the p53-dependent DNA damage response pathway by binding to and deacetylating p53, specifically at Lysine 382.

Storage


Storage: Store at -80°C. Storage Buffer:

PBS Only

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82790-4-RR (SIRT1 antibody) at dilution of 1:4700 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SIRT1 antibody (82790-4-RR, Clone: 2F11) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004). This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.