

For Research Use Only

# Mono-Methyl-Histone H3 (Lys9) Recombinant antibody

Catalog Number: 82821-6-RR



## Basic Information

<b>Catalog Number:</b> 82821-6-RR	<b>GenBank Accession Number:</b> BC066245	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ul , Concentration: 1000 ug/ml by Nanodrop;	<b>GeneID (NCBI):</b> 8350	<b>CloneNo.:</b> 3C23
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P68431	<b>Recommended Dilutions:</b> WB 1:5000-1:50000
<b>Isotype:</b> IgG	<b>Full Name:</b> histone cluster 1, H3a	
	<b>Observed MW:</b> 15 kDa	

## Applications

<b>Tested Applications:</b> WB, Dot Blot, ELISA	<b>Positive Controls:</b>
<b>Species Specificity:</b> human, rat	<b>WB :</b> HeLa cells, HEK-293 cells, K-562 cells, THP-1 cells, HSC-T6 cells

## Background Information

Histones, including H1/H5 (linker histones), H2, H3, and H4 (core histones), are nucleic proteins which interact with DNA to form the nucleosomes and play important roles in gene regulation and DNA replication. Histone proteins are highly post-translationally modified while Histone H3 is the most extensively modified. Methylation of Histone H3 at lysine 9 is linked to transcriptional repression.

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**

\*\*\* 20ul sizes contain 0.1% BSA



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Dot blot analysis was used to confirm the specificity of 82821-6-RR Mono-Methyl-Histone H3 (Lys9) antibody. Peptides were spotted onto NC and probed with antibody at 1 µg/ml. The amount of peptide (ug/mL) spotted is indicated next to each row.

Various lysates were subjected to SDS PAGE followed by western blot with 82821-6-RR (Mono-Methyl-Histone H3 (Lys9) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.