For Research Use Only

Acetyl-Histone H2A (Lys9) Recombinant antibody, PBS Only proteintech® Antibodies | ELISA kits | Proteins Uni-rAb www.ptglab.com

Catalog Number:82823-2-PBS

Basic Information

Catalog Number: 82823-2-PBS Size: 100ug , Concentration: 1 mg/ml by Nanodrop; Source: Rabbit Isotype: IgG

GenBank Accession Number: BC062211 GeneID (NCBI): 221613 UNIPROT ID: Q96QV6 Full Name: histone cluster 1, H2aa Observed MW: 16 kDa

Purification Method: Protein A purfication CloneNo.: 1017

Applications

Tested Applications: WB, Dot Blot, Indirect ELISA Species Specificity: human

Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data

Peptide Conc. (ug/ml)	1	2	3	4	5	6	7	8	9				
(ug/mi) 12.5												100 kDa→	1
	-											70 kDa→	
1.25												50 kDa→	
												40 kDa→	
0.125												H	
0.0125												30 kDa-	
0.0125												5	
0.00125												20 kDa→	
												Ö	
			Antil	oody C	onc: 1	ıg/mL						0	
Dot blot analysis wa												15 kDa→	
were spotted onto N indicated next to eac		ied with	antibot	iy at 1 µ	g/mi.in	e amoun	it of pept	ide (µg/n	it) spotte				
Column 1: H2AK9ac.													
Column 5: H2AK13A H2AK15, Column 9: I			dified H	IZAK13.	Column	7: H2AK	15ac. Col	umn 8:U	nmodifie	d			

Ŧ

Dot blot analysis was used to confirm the specificity of Acetyl-Histone H2A (Lys9) antibody. Acetylated peptides were spotted onto NC and probed with antibody at 1 µg/ml.The amount of peptide (µg/mL) spotted is indicated next to each row. Column 1: H2AK9ac. Column 2: Unmodified H2AK9. Column 3: H2AK5ac. Column 4: Unmodified H2AK5. Column 5: H2AK13Ac. Column 6: Unmodified H2AK13. Column 7: H2AK15ac. Column 8:Unmodified H2AK15. Column 9: Column 8: Unmodified H2AK15. Column 9:

(66031-1-lg) mi 4h

Trichostain A treated and untreated HeLa cell lysates were subjected to SDS PAGE followed by western blot with 82823-2-RR (Acetyl-Histone H2A (Lys9) antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82823-2-PBS in a different storage buffer formulation formulation.