For Research Use Only

CD147 Recombinant antibody

Catalog Number:82909-7-RR




Basic Information	Catalog Number: 82909-7-RR	GenBank Accession Number: BC 009040	Purification Method: Protein A purfication
	Size: 100ul , Concentration: 1000 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG	GeneID (NCBI): 682	CloneNo.: 230046A9
		UNIPROT ID: P35613 Full Name: basigin (Ok blood group)	Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:500-1:2000
		Observed MW: 50-60 kDa	
		Applications	Tested Applications: WB, IF/ICC, ELISA Species Specificity: human
WB : HeLa cells, HEK-293T cells IF/ICC : HeLa cells,			
Background Information	CD147, also known as Basigin or extracellular matrix metalloproteinase inducer (EMMPRIN), is a transmembrane glycoprotein that belongs to the immunoglobulin superfamily (PMID: 7812975). The molecule is composed of an intracellular portion, an extracellular portion and a single transmembrane region. CD147 is expressed on a variety of cell types (e.g., hematopoietic, epithelial, and endothelial cells) and at varying levels (PMID: 32968061). Increased expression of CD147 occurs in many tumors. CD147 is a pleiotropic molecule that plays an important role in fetal, neuronal, lymphocyte and extracellular matrix development (PMID: 17945211). CD147 has been identified as a receptor essential for erythrocyte invasion by Plasmodium falciparum (PMID: 22080952). It has been reported that spike protein of SARS-CoV-2 binds to CD147 on host cells, thereby mediating the viral invasion (Wang, Ke, et al, BioRxiv, 2020).		
Storage	Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°		
	Aliquoting is unnecessary for -20°C st	torage	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




WWW.PTGLAB.COM

Various lysates were subjected to SDS PAGE followed by western blot with 82909-7-RR (CD147 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CD147 antibody (82909-7-RR, Clone: 230046A9) at dilution of 1:1000 and MultirAb CoraLite ® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004).

Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CD147 antibody (82909-7-RR, Clone: 230046A9) at dilution of 1:200 and MultirAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).



Biolayer interferometry (BLL) kinetic assays of 82909-7-RR against Human CD147 were performed. The affinity constant is 21.7 nM.