

For Research Use Only

CYP7B1 Recombinant antibody

Catalog Number: 82927-1-RR



Basic Information

Catalog Number: 82927-1-RR	GenBank Accession Number: BC136574	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 9420	CloneNo.: 230160H5
Source: Rabbit	UNIPROT ID: O75881	Recommended Dilutions: WB 1:2000-1:10000 IF/ICC 1:500-1:2000
Isotype: IgG	Full Name: cytochrome P450, family 7, subfamily B, polypeptide 1	
Immunogen Catalog Number: AG19374	Calculated MW: 506 aa, 58 kDa	
	Observed MW: 50-55 kDa	

Applications

Tested Applications: WB, IF/ICC, FC (Intra), ELISA	Positive Controls: WB : A549 cells, HepG2 cells, HeLa cells, HCT 116 cells IF/ICC : HepG2 cells, HEK-293 cells
Species Specificity: human	

Background Information

CYP7B1(Cytochrome P450 7B1) is also named as 25-hydroxycholesterol 7-alpha-hydroxylase and belongs to the cytochrome P450 family. It catalyzes the first and rate-limiting step in the coconversion of cholesterol to bile acids and CYP7B1-mediated catalysis may play a role for control of the cellular levels of androgens, not only of estrogens.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82927-1-RR (CYP7B1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using CYP7B1 antibody (82927-1-RR, Clone: 230160H5) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



1×10^6 HeLa cells were intracellularly stained with 0.25 μ g CYP7B1 Recombinant antibody (82927-1-RR, Clone:230160H5) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 μ g Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).