For Research Use Only

Integrin beta 3 Recombinant antibody

83053-4-RR

Catalog Number:83053-4-RR



Basic Information

Catalog Number: GenBank Accession Number:

Purification Method: BC127667 Protein A purification

GeneID (NCBI): CloneNo.: 100ul, Concentration: 1000 ug/ml by 3690 230292D5

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source: P05106 IF/ICC 1:200-1:800

Rabbit Full Name:

Isotype: integrin, beta 3 (platelet glycoprotein

IIIa, antigen CD61) IgG

> Calculated MW: 87 kDa Observed MW: 100-110 kDa

Applications

Tested Applications: IF/ICC, FC (Intra), ELISA

Species Specificity:

Positive Controls:

IF/ICC: HUVEC cells,

Background Information

Integrin beta-3, also named as CD61 and GPIIIa, is a receptor for cytotactin, fibronectin, laminin, matrix metallo protein a se-2, osteopontin, osteomodulin, prothrombin, thrombos pondin, vitronectin and von Willebrand and von Willefactor. Integrin beta-3 recognize the sequence R-G-D in a wide array of ligands and the sequence H-H-L-G-G-G-A-K-Q-A-G-D-V in fibrinogen gamma chain. Following activation integrin beta-3 brings about platelet/platelet interaction through binding of soluble fibrinogen which leads to rapid platelet aggregation which physically plugs ruptured endothelial surface. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance $angiogenesis in Kaposi's sarcoma\ lesions.\ Human\ integrin\ beta-3\ has\ a\ calculated\ molecular\ weight\ of\ 87\ kDa.\ As\ a$ result of glycosylation, the apparent molecular mass of integrin beta-3 is approximately 90-110 kDa.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HUVEC cells using Integrin beta-3 antibody (83053-4-RR, Clone: 230292D5) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10^6 HUVEC cells were intracellularly stained with 0.25 ug Anti-Human Integrin beta-3 (83053-4-RR, Clone:230292D5) and APC-Conjugated Goat Anti-Rabbit 1gG(H+L)(red), or 0.25 ug Rabbit 1gG control Rabbit PolyAb (3000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).