## For Research Use Only

## STK39 Recombinant antibody

Catalog Number:83081-2-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

 83081-2-RR
 BC166614

 Size:
 GeneID (NCBI):

 100ul , Concentration: 1000 µg/ml by 27347

Nanodrop; UNIPROT ID:
Source: Q9UEW8
Rabbit Full Name:

Isotype: serine threonine kinase 39
IgG (STE20/SPS1 homolog, yeast)

Immunogen Catalog Number: Calculated MW: AG34503 59 kDa

Observed MW: 62 kDa

**Applications** 

Tested Applications:

WB, IF/ICC, IF-P, ELISA

Species Specificity: human, mouse, rat

**Positive Controls:** 

WB: Jurkat cells, MCF-7 cells, SH-SY5Y cells, HEK-293T

**Purification Method:** 

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000

IF/ICC 1:250-1:1000

IF-P 1:50-1:500

CloneNo.:

230382G9

cells, HepG2 cells, rat brain tissue

IF-P: mouse brain tissue,
IF/ICC: NIH/3T3 cells,

## **Background Information**

STK39 (Serine/Threonine Kinase 39) is a proline- and alanine-rich Ste20-related kinase (also named as SPAK), which is composed of a short N-terminal proline and alanine repeats (PAPA box), a kinase catalytic domain and a C-terminal regulatory domain (PMID: 16382158). In mammalians, STK39 plays an important role in ion homeostasis by regulating the cation chloride cotransporters' activities, which is crucial for the modulation of renal salt transport and blood pressure (PMID: 33500714). Notably, STK39 is also implicated in the regulation of tumor cell proliferation, migration, and invasion in multiple cancers, including osteosarcoma and cervical cancer (PMID: 34335956).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 83081-2-RR (STK39 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using STK39 antibody (83081-2-RR, Clone: 230382G9) at dilution of 1:500 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using STK39 antibody (83081-2-RR, Clone: 230382G9) at dilution of 1:200 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9 0)



Biolayer interferometry (BLL) kinetic assays of 83081-2-RR against Human STK39 were performed. The affinity constant is 2.52 nM.