

For Research Use Only

NLRP10 Recombinant antibody

Catalog Number: 83131-8-RR



Basic Information

Catalog Number: 83131-8-RR	GenBank Accession Number: BC104957	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 338322	CloneNo.: 230402C1
Source: Rabbit	UNIPROT ID: Q86W26	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: NLR family, pyrin domain containing 10	
Immunogen Catalog Number: AG26402	Calculated MW: 655 aa, 75 kDa	
	Observed MW: 75 kDa	

Applications

Tested Applications:
WB, FC (Intra), ELISA

Species Specificity:
human, mouse

Positive Controls:
WB : mouse heart tissue,

Background Information

NLRP10 is a member of the NALP protein family typically contain a NACHT domain, a NACHT-associated domain (NAD), a C-terminal leucine-rich repeat (LRR) region, and an N-terminal pyrin domain (PYD). NLRP10 likely plays a regulatory role in the innate immune system. The protein belongs to the signal-induced multiprotein complex, the inflammasome, that activates the pro-inflammatory caspases, caspase-1 and caspase-5.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.

Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



mouse heart tissue were subjected to SDS PAGE followed by western blot with 83131-8-RR (NLRP10 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1×10^6 A431 cells were intracellularly stained with 0.25 ug Nlrp10 Recombinant Antibody (83131-8-RR, Clone:230402C1) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.



Biolayer interferometry (BLI) kinetic assays of 83131-8-RR against Human NLRP10 were performed. The affinity constant is 0.899 nM.