For Research Use Only

TLR3 Recombinant antibody

Catalog Number:83136-3-RR 1 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purfication

83136-3-RR

GeneID (NCBI):

Size:

CloneNo.:

IHC 1:200-1:800

100ul , Concentration: 700 $\mu g/ml$ by Nanodrop;

BC094737

230445G4

UNIPROT ID: 015455

Recommended Dilutions: WB 1:2000-1:14000

Rabbit

Full Name:

toll-like receptor 3

Isotype: IgG

Calculated MW:

Immunogen Catalog Number: AG34044

904 aa, 104 kDa

Observed MW:

120 kDa, 68 kDa

Applications

Tested Applications:

WB, IHC, ELISA

WB: HT-29 cells, human placenta tissue

IHC: human ovarian cancer, human placenta tissue

Positive Controls:

Cited Applications: WB

Species Specificity:

human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

TLR3 belongs to the Toll-like receptor family which is important in the innate immune response to pathogens. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. TLR3 is a nucleotide-sensing TLR that is activated by double-stranded RNA. Upon recognition of dsRNA, TLR3 transmits signals via the adaptor protein TICAM-1/TRIF, leading to the induction of type I IFN, cytokine/chemokine production, and dendritic cell (DC) maturation (PMID: 18262679).

Notable Publications

Author	Pubmed ID	Journal	Application
Tamaki Suganuma	39465205	NAR Mol Med	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data




HT-29 cells were subjected to SDS PAGE followed by western blot with 83136-3-RR (TLR3 antibody) at dilution of 1:7000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human ovarian cancer slide using 83136-3-RR (TLR3 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovarian cancer slide using 83136-3-RR (TLR3 antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLL) kinetic assays of 83136-3-RR against Human TLR3 were performed. The affinity constant is below 1 pM.