For Research Use Only

SMARCA4 Recombinant antibody

Catalog Number:83310-7-RR



Basic Information

Catalog Number: GenBank Accession Number:

83310-7-RR BC150298 Protein A purfication
Size: GeneID (NCBI): CloneNo.:

100ul , Concentration: 1000 µg/ml by 6597 240172F7

 Nanodrop;
 UNIPROT ID:
 Recommended Dilutions:

 Source:
 P51532
 WB 1:5000-1:50000

 Rabbit
 Full Name:
 IHC 1:50-1:500

Isotype: SWI/SNF related, matrix associated, IgG actin dependent regulator of

Immunogen Catalog Number: chromatin, subfamily a, member 4

AG16256 Calculated MW:

1647 aa, 185 kDa Observed MW: 185 kDa

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, HepG2 cells, HSC-T6 cells, NIH/3T3

Purification Method:

cells, HEK-293T cells, C6 cells

IHC: human colon cancer tissue, mouse kidney tissue

Background Information

SMARCA4, also named as BAF190A, BRG1, SNF2B and SNF2L4, belongs to the SNF2/RAD54 helicase family. SMARCA4 is a transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. It is a component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator complex. It is also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene.

Storage

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83310-7-RR (SMARCA4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 83310-7-RR (SMARCA4 antibody) at dilution of 1:10000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 83310-7-RR (SMARCA4 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 83310-7-RR (SMARCA4 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 Jurkat cells were intracellularly stained with 0.25 ug SMARCA4 Recombinant antibody (83310-7-RR, Clone:240172F7) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLL) kinetic assays of 83310-7-RR against Human SMARCA4 were performed. The affinity constant is 1.73 nM.