

For Research Use Only

MUC1/CA15-3 C-terminal Recombinant antibody, PBS Only

Catalog Number: 83311-4-PBS



Basic Information

Catalog Number: 83311-4-PBS	GenBank Accession Number: BC120975	Purification Method: Protein A purification
Size: 100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 4582	CloneNo.: 240190G10
Source: Rabbit	UNIPROT ID: P15941	
Isotype: IgG	Full Name: mucin 1, cell surface associated	
Immunogen Catalog Number: AG20366	Calculated MW: 1264 aa, 123 kDa	
	Observed MW: 23-30 kDa	

Applications

Tested Applications:
WB, IHC, ELISA

Species Specificity:
human, mouse

Background Information

MUC1 is a type I transmembrane glycoprotein expressed by various epithelial cells of the female reproductive tract, lung, breast, kidney, stomach, and pancreas. MUC1 is transcribed as a large precursor gene product, and upon translation, is cleaved in the endoplasmic reticulum, yielding two subunits: the large extracellular N-terminal subunit (MUC1-N, about 120-200 kDa) and the small cytoplasmic C-terminal subunit (MUC1-C, about 23-30 kDa). Among the known mucins, MUC1 is best studied and plays crucial roles in regulating many cellular properties, including cell proliferation, apoptosis, adhesion, and invasion. MUC1 is overexpressed in a wide range of human epithelial malignancies. This antibody was raised against the C-terminal region of human MUC1, thus it recognizes the 23-30 kDa cytoplasmic MUC1.

Storage


Storage:
Store at -80°C.

Storage Buffer:
PBS Only


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83311-4-RR (MUC1/CA15-3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83311-4-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 83311-4-RR (MUC1/CA15-3 C-terminal antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83311-4-PBS in a different storage buffer formulation.


Immunohistochemical analysis of paraffin-embedded human rectal cancer tissue slide using 83311-4-RR (MUC1/CA15-3 C-terminal antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83311-4-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 83311-4-RR (MUC1/CA15-3 C-terminal antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83311-4-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 83311-4-RR (MUC1/CA15-3 C-terminal antibody) at dilution of 1:800 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83311-4-PBS in a different storage buffer formulation.



Bi-layer interferometry (BLI) kinetic assays of 83311-4-RR against Human CA15-3/MUC1 were performed. The affinity constant is 63.7 pM.