For Research Use Only

NCAM1/CD56 Recombinant antibody, PBS Only (Capture)

Catalog Number:83365-5-PBS



Basic Information

Catalog Number: 83365-5-PBS

BC047244

Purification Method: Protein A purfication

Size:

Source

Rabbit

Isotype

IgG

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by

CloneNo.: 240078A7

Nanodrop:

ENSEMBL Gene ID: ENSG00000149294 **UNIPROT ID:**

P13591 Full Name:

neural cell adhesion molecule 1

GenBank Accession Number:

Calculated MW: 95 kDa

Applications

Tested Applications:

Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity:

human

Product Information

83365-5-PBS targets NCAM1/CD56 as part of a matched antibody pair:

MP00393-4: 83365-5-PBS capture and 83365-4-PBS detection (validated in Sandwich ELISA)

Unconjugated rabbit recombinant monoclonal antibody in PBS only (BSA and azide free) storage buffer at a $concentration of 1\,mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant$ technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.


Background Information


Neural cell adhesion molecule 1 (NCAM1, also known as CD56) is a cell adhesion glycoprotein of the $immuno globulin (Ig) \, superfamily. \, It \, is \, a \, multifunction \, protein \, involved \, in \, synaptic \, plasticity, \, neurodevelopment, \, in \, involved \, in \, synaptic \, plasticity, \, neurodevelopment, \, in \, involved \, in \, synaptic \, plasticity, \, neurodevelopment, \, involved \, in \, synaptic \, plasticity, \, neurodevelopment, \, involved \, in \, synaptic \, plasticity, \, neurodevelopment, \, involved \, i$ and neurogenesis. NCAM1 is expressed on human neurons, glial cells, skeletal muscle cells, NK cells and a subset of T cells, and the expression is observed in a wide variety of human tumors, including myeloma, myeloid leukemia, neuroendocrine tumors, Wilms' tumor, neuroblastoma, and NK/T cell lymphomas. Three major isoforms of NCAM1, with molecular masses of 120, 140, and 180 kDa, are generated by alternative splicing of mRNA (PMID: 9696812). The glycosylphosphatidylinositol (GPI)-anchored NCAM120 and the transmembrane NCAM140 and NCAM180 consist of five Ig-like domains and two fibronection-type III repeats (FNIII). All three forms can be posttranslationally modified by addition of polysialic acid (PSA) (PMID: 14976519). Several other isofroms have also been described (PMID: 1856291).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data







The NCAM1/CD56 concentration of Human cerebrospinal fluid (CSF) samples were determined to be 127.3 ng/mL



Serum of sixteen individual healthy human donors were measured. The NCAM1/CD56 concentration of detected samples was determined to be 1.2 ug/mL with a range of 0.7 - 1.9 ug/mL