For Research Use Only

SIAH1 Recombinant antibody

Catalog Number:83389-4-RR



Basic Information

Catalog Number: GenBank Accession Number:

83389-4-RR BC035562 GeneID (NCBI): Size:

100ul , Concentration: 1000 $\mu g/ml$ by 6477

Nanodrop: **UNIPROT ID:** Q8IUQ4 Rabbit Full Name:

Isotype: seven in absentia homolog 1

(Drosophila) IgG Immunogen Catalog Number: Calculated MW: 282 aa, 31 kDa AG34618 Observed MW:

30-34 kDa

Applications

Tested Applications:

WB, IHC, ELISA Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, mouse kidney tissue, HepG2 cells, HEK-293T cells, SH-SY5Y cells, Jurkat cells, COLO 320

Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:50-1:500

CloneNo.:

240377B4

IHC: mouse testis tissue,

Background Information

SIAH1, also named as HUMSIAH, Siah-1a and Siah-1S(21kd), belongs to the SINA (Seven in absentia) family. SIAH1 is known to cause indirect degradation of beta-catenin through formation of a complex with Siah-interacting protein (SIP), Skp1 and Ebi. SIAH1 can form heterodimer or homodimer such as Siah-1*Siah-1(70 or 62kd), Siah-1*Siah-1 1S(42kd) and Siah-1S*Siah-1S(51-56kd). Importantly, results from in vitro soft agar assay demonstrated that Siah-1S displays a promotion effect on cells tumorigenicity. (PMID:17420721)

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83389-4-RR (SIAH1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




mouse kidney tissue were subjected to SDS PAGE followed by western blot with 83389-4-RR (SIAH1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 83389-4-RR (SIAH1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 A549 cells were intracellularly stained with 0.25 ug Siah1 Recombinant Antibody (83389-4-RR, Clone:240377B4) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83389-4-RR against Human SIAH1 were performed. The affinity constant is 2.39 nM.