

For Research Use Only

NCAM2 Recombinant antibody

Catalog Number: 83494-2-RR



Basic Information

Catalog Number: 83494-2-RR	GenBank Accession Number: BC052946	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 4685	CloneNo.: 240398C5
Source: Rabbit	UNIPROT ID: O15394	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: neural cell adhesion molecule 2	
Immunogen Catalog Number: AG4821	Calculated MW: 93 kDa	
	Observed MW: 90 kDa, 125 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : mouse brain tissue, mouse cerebellum tissue, rat brain tissue
Species Specificity: Human, mouse , rat	

Background Information

The mammalian NCAM family has two members, NCAM1 and NCAM2 (PMID: 9729879; 20439718). Neural cell adhesion molecule 2 (NCAM2, also known as NCAM21) is a single-pass type I membrane protein that is highly expressed in the brain, which may play important roles in selective fasciculation and zone-to-zone projection of the primary olfactory axons or in neurodegenerative diseases such as Alzheimer's disease (PMID: 26611261). NCAM2 has two isoforms: NCAM2.1 bearing transmembrane and cytoplasmic domains and the shorter NCAM2.2 isoform, which is GPI-anchored (PMID: 9334170; 9221781). The molecular weight of NCAM2 is around 125 kDa and 93 kDa (PMID: 9334170; 3171685).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83494-2-RR (NCAM2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 83494-2-RR against Human NCAM2 were performed. The affinity constant is 0.882 nM.