

For Research Use Only

TCF7L1 Recombinant antibody

Catalog Number:83588-3-RR



Basic Information

Catalog Number: 83588-3-RR	GenBank Accession Number: BC058894	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 83439	CloneNo.: 240544F1
Source: Rabbit	UNIPROT ID: Q9HCS4	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG	Full Name: transcription factor 7-like 1 (T-cell specific, HMG-box)	
Immunogen Catalog Number: AG34253	Calculated MW: 63 kDa	
	Observed MW: 63 kDa, 75 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : A431 cells, HeLa cells, HEK-293 cells, NCCIT cells, NIH/3T3 cells
Species Specificity: human, mouse	

Background Information

TCF7L1 is one of TCF/LEF transcription factors that are mediators of the Wnt signaling pathway and are antagonized by the TGF-beta signaling pathway. It binds to DNA and acts as a repressor in the absence of CTNNB1, and as an activator in its presence. It's necessary for the terminal differentiation of epidermal cells, the formation of keratohyalin granules and the development of the barrier function of the epidermis

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83588-3-RR (TCF7L1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



1×10^6 A431 cells were intracellularly stained with 0.25 μ g TCF7L1 Recombinant antibody (83588-3-RR, Clone:240544F1) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 μ g Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).