

For Research Use Only

Flotillin 1 Recombinant antibody

Catalog Number: 83598-4-RR



Basic Information

Catalog Number: 83598-4-RR	GenBank Accession Number: BC001146	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 ug/ml by Nanodrop;	GeneID (NCBI): 10211	CloneNo.: 240582G6
Source: Rabbit	UNIPROT ID: O75955	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: flotillin 1	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
Immunogen Catalog Number: AG7902	Calculated MW: 47 kDa	IHC 1:200-1:800
	Observed MW: 47 kDa	IF/ICC 1:250-1:1000

Applications

Tested Applications:
WB, IHC, IF/ICC, IP, ELISA

Species Specificity:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : U-87 MG cells, LNCaP cells, Raji cells, mouse heart tissue, mouse brain tissue, rat brain tissue

IP : mouse brain tissue,

IHC : human stomach cancer tissue,

IF/ICC : HeLa cells,

Background Information

Flotillins are membrane-associated proteins that contain two homologous isoforms, flotillin 1 (FLOT1) and flotillin 2 (FLOT2). They are highly conserved and widely expressed proteins commonly used as lipid raft markers. Flotillins are involved in various signaling pathways, cell adhesion, membrane trafficking and axonal growth.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83598-4-RR (Flotillin 1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 83598-4-RR (Flotillin 1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 83598-4-RR (Flotillin 1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-Flotillin 1 (IP:83598-4-RR, 4ug; Detection:83598-4-RR 1:3000) with mouse brain tissue lysate 2400 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Flotillin 1 antibody (83598-4-RR, Clone: 240582G6) at dilution of 1:500 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).