For Research Use Only

PD-L1/CD274 Recombinant antibody, PBS Only (Capture)

Catalog Number:83600-7-PBS



Basic Information

Catalog Number:

GenBank Accession Number:

BC074984

Purification Method: Protein A purification

CloneNo.:

240721D4

83600-7-PBS Size:

GeneID (NCBI):

100ug, Concentration: 1 mg/ml by

29126

UNIPROT ID:

Source: Rabbit

Nanodrop:

Q9NZQ7

Isotype:

IgG

Full Name: CD274 molecule

Calculated MW: 290 aa, 33 kDa

Applications

Tested Applications:

Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity:

human

Product Information

83600-7-PBS targets PD-L1/CD274 as part of a matched antibody pair:

MP00601-4: 83600-7-PBS capture and 83600-6-PBS detection (validated in Sandwich ELISA)


Unconjugated rabbit recombinant monoclonal antibody in PBS only (BSA and azide free) storage buffer at a $concentration of 1\,mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant$ technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data




Sandwich ELISA standard curve of MP00601-4, Human PD-L1/CD274 Recombinant Matched Antibody Pair - PBS only. 83600-7-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Eg1114. 83600-6-PBS was HRP conjugated as the detection antibody. Range: 3.9-250 pg/mL




Human peripheral blood mononuclear cells (PBMC) were cultured unstimulated or stimulated with 10 µg/mL PHA for 3 days or 5-days. For the PBMC supernatant cultured for 3 days, the mean PD-L1/CD274 concentration was determined to be 7.7 pg/mL in unstimulated PBMC supernatant, 108.4 pg/mL in PHA stimulated PBMC supernatant. For the PBMC supernatant cultured for 5 days, the mean PD-L1/CD274 concentration was determined to be 11.0 pg/mL in



Urine of seven individual healthy human donors was measured. The PD-L1/CD274 concentration of detected samples was determined to be 10.8 pg/mL with a range of 0.7 - 33.1 pg/mL



The mean PD-L1/CD274 concentration was determined to be 36,368.3 pg/mL in MDA-MB-231 cell extract based on a 2.3 mg/mL extract load.



Serum of sixteen individual healthy human donors was measured. The PD-L1/CD274 concentration of detected samples was determined to be 75.1 pg/mL with a range of 34.4-168.1 pg/mL.