For Research Use Only

ASGR1 Recombinant antibody

Catalog Number:83691-3-RR



Basic Information

Catalog Number: GenBank Accession Number:

83691-3-RR NM_001671.4 GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 432 Nanodrop: **UNIPROT ID:**

Source: P07306-1 Rabbit Full Name:

Isotype: asialoglycoprotein receptor 1 IgG

Calculated MW:

33 kDa Observed MW: 42-46 kDa

Purification Method:

Protein A purification

CloneNo.: 240672C4

Recommended Dilutions: WB 1:1000-1:4000 IHC 1:500-1:2000 IF-P 1:200-1:800

Applications

Tested Applications: WB, IHC, IF-P, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HepG2 cells, HuH-7 cells, human liver tissue, rat

liver tissue

IHC: mouse liver tissue, IF-P: rat liver tissue,

Background Information

Asialoglycoprotein receptor (ASGPR), also known as the hepatic galactose/N-acetylglucosamine (GlcNAc) receptor or Ashwell receptor, is a C-type lectin expressed exclusively in hepatic parenchymal cells. ASGPR consists of two subunits, a major subunit (ASGR1, HL-1) and a minor subunit (ASGR2, HL-2), and specifically recognizes terminal βlinked galactose or GlcNAc on circulating glycoproteins or cells. This receptor plays a critical role in serum glycoprotein homeostasis by mediating the endocytosis and lysosomal degradation of glycoproteins that contain terminal galactose or GlcNAc residues. ASGPR may facilitate hepatic infection by multiple viruses including hepatitis B, and is also a target for liver-specific drug delivery.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83691-3-RR (ASGR1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 83691-3-RR (ASGR1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 83691-3-RR (ASGR1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat liver tissue using ASGR1 antibody (83691-3-RR, Clone: 240672C4) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat liver tissue using ASGR1 antibody (83691-3-RR, Clone: 240672C4) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLL) kinetic assays of 83691-3-RR against Human ASGR1 were performed. The affinity constant is 1.47 nM.