

For Research Use Only

# RXRA Recombinant antibody, PBS Only

Catalog Number: 83796-5-PBS



## Basic Information

<b>Catalog Number:</b> 83796-5-PBS	<b>GenBank Accession Number:</b> BC110998	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ug, Concentration: 1 mg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 6256	<b>CloneNo.:</b> 240878A9
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P19793	
<b>Isotype:</b> IgG	<b>Full Name:</b> retinoid X receptor, alpha	
<b>Immunogen Catalog Number:</b> AG15651	<b>Calculated MW:</b> 462 aa, 51 kDa	
	<b>Observed MW:</b> 54 kDa, 44 kDa	

## Applications

**Tested Applications:**  
WB, IF/ICC, FC (Intra), ELISA

**Species Specificity:**  
human, mouse

## Background Information

Retinoid X receptor alpha (RXRA). Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. The RAR/RXR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. The high-affinity ligand for RXRs is 9-cis retinoic acid. RXRA serves as a common heterodimeric partner for a number of nuclear receptors. The RXR/RAR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. In the absence of a ligand, the RXR-RAR heterodimers associate with a multiprotein complex containing transcription corepressors that induce histone acetylation, chromatin condensation, and transcriptional suppression. On ligand binding, the corepressors dissociate from the receptors and associate with the coactivators leading to transcriptional activation. The RXRA/PPARA heterodimer is required for PPARA transcriptional activity on fatty acid oxidation genes such as ACOX1 and the P450 system genes. This antibody is a rabbit polyclonal antibody raised against the 350 AA of human RXRA C-terminal. RXRA is highly expressed in the liver and also expressed in the lungs, kidneys, and heart. It can recognize the mature 54 kDa RXRA and the truncated 44 kDa RXRA (PMID: 20541701).

## Storage


**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83796-5-RR (RXRA antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83796-5-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (4% PFA) fixed HeLa cells using RXRA antibody (83796-5-RR, Clone: 240878A9) at dilution of 1:600 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 83796-5-PBS in a different storage buffer formulation.



$1 \times 10^6$  MCF-7 cells were intracellularly stained with 0.25  $\mu$ g RXRA Recombinant antibody (83796-5-RR, Clone:240878A9) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25  $\mu$ g Isotype Control (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set. This data was developed using the same antibody clone with 83796-5-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 83796-5-RR against Human RXRA were performed. The affinity constant is below 1 pM.