

For Research Use Only

Beta Arrestin 1 Recombinant antibody

Catalog Number:83837-1-RR



Basic Information

Catalog Number: 83837-1-RR	GenBank Accession Number: BC003636	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 408	CloneNo.: 240974C3
Source: Rabbit	UNIPROT ID: P49407	Recommended Dilutions: WB 1:1000-1:8000
Isotype: IgG	Full Name: arrestin, beta 1	
Immunogen Catalog Number: AG7608	Calculated MW: 47 kDa	
	Observed MW: 47-55 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HEK-293 cells, RAW 264.7 cells, mouse brain tissue, rat lung tissue
Species Specificity: human, mouse, rat	

Background Information

β-Arrestins (ARRBs), the best known regulators of G protein-coupled receptor signaling, are versatile and multifunctional adapter proteins that regulate diverse cellular functions, including cell growth, apoptosis and immune responses. Overexpression of beta Arrestin 1 has been found in various cancers, indicating it as a potential therapeutic target for cancer treatment. Recently expression of ARRB1 in saliva has been identified as a candidate circadian biomarker. ARRB1 migrated as a doublet of two bands of 45 and 55 kDa (PMID:28947386).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83837-1-RR (Beta Arrestin 1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 83837-1-RR (Arrestin antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 83837-1-RR against Human Beta Arrestin 1 were performed. The affinity constant is 0.759 nM.