For Research Use Only

NUCB2/nesfatin-1 Recombinant antibody, PBS Only

Catalog Number:83917-5-PBS



Basic Information

Catalog Number: 83917-5-PBS

GenBank Accession Number:

Purification Method: NM_005013 Protein A purfication

CloneNo.:

241037C1

GeneID (NCBI): Size: 100ug, Concentration: 1 mg/ml by

Nanodrop: **UNIPROT ID:** P80303 Rabbit Full Name: Isotype: nucleobindin 2 IgG

Calculated MW:

Immunogen Catalog Number: 50 kDa

AG25152

Tested Applications:

IF/ICC, FC (Intra), Indirect ELISA

Species Specificity:

Background Information


NucB2, an anorexigenic molecule, is expressed mainly in the hypothalamus, particularly in the supraoptic nucleus (SON) and the paraventricular nucleus (PVN). NucB2 is also expressed in the subfornical organ (SFO). Because the $SON \ and \ PVN \ are involved \ in \ body \ fluid \ regulation, \ Nuc B2 \ may \ be involved \ in \ dehydration-induced$ anorexia. NUCB2 is composed of a signal peptide of 24 amino acids in the N-terminal region and a protin structure containing 396 amino acids. Nesfatin-1 is an 82-amino-acid peptide hormone cleaved from nucleobindin2 (NUCB2).

Storage


Applications

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using NUCB2 antibody (83917-5-RR, Clone: 241037C1) at dilution of 1:250 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) (SA00013-2). This data was developed using the same antibody clone with 83917-5-PBS in a different storage buffer formulation.



1x10^6 A431 cells were intracellularly stained with 0.25 ug NUCB2 Recombinant antibody (83917-5-RR, Clone:241037C1) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 83917-5-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using NUCB2/nesfatin-1 antibody (83917-5-RR, Clone: 241037C1) at dilution of 1:500 and CoraLite® 594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4). This data was developed using the same antibody clone with 83917-5-PBS in a different storage buffer formulation.