For Research Use Only

Phospho-AMPK Beta 1 (Ser182) Recombinant antibody, PBS Only

Catalog Number:83924-1-PBS




Basic Information	Catalog Number: 83924-1-PBS	GenBank Accession Number: BC001007	Purification Method: Protein A purfication
	Size:	GenelD (NCBI):	CloneNo.:
	100ug , Concentration: 1 mg/ml by	5564	240628B1
	Nanodrop;	UNIPROT ID:	
	Source:	Q9Y478	
	Rabbit	Full Name:	
	Isotype: IgG	protein kinase, AMP-activated, beta 1 non-catalytic subunit	
	-	Calculated MW: 38 kDa	
		Observed MW:	
		38 kDa	
Applications	Tested Applications: WB, ELISA		
	Species Specificity:		
	human		
Background Information	AMPK Beta 1 (5'-AMP-activated prote	ein kinase subunit beta-1) is also na	med as PRKAB1 and AMPK. AMPK, a
background mormation	serine/threonine kinase that exists as a heterotrimer comprised of a catalytic α-subunit and regulatory β- and γ- subunits, has been recognized as a sensor of cellular energy homeostasis (PMID: 21937710). AMPK regulates key metabolic enzymes, cell growth, apoptosis, gene transcription, and protein synthesis (PMID: 12829246). AMPK is an energy sensor and plays an essential role in the control of cellular bioenergetics by responding to various stresses including those that induce changes in the cellular AMP:ATP ratio or modulation in intracellular calcium (PMID: 27812976, PMID: 26616193). Recent studies have shown that AMPK mediates the inhibition of cell proliferation and growth of tumor cells (PMID: 16613876). AMPK also inhibits the expression of Glut1 and glycolysis in Tregs by inhibiting mTORC1 signaling (PMID: 25477880). This antibody recognizes phosphorylated AMPK Beta 1.		
Storage	Storage: Store at -80°C. Storage Buffer: PBS Only		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



٠

•

Non-treated HEK-293 cells, phosphatase inhibitor treated HEK-293 cells and λ phosphatase treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 83924-1-RR (Phospho-AMPK Beta 1 (Ser182) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-lg) antibody as a loading control. This data was developed using the same antibody clone with 83924-1-