For Research Use Only

EOMES/TBR2 Recombinant antibody

Catalog Number:83945-5-RR



Purification Method:

Protein A purfication

Basic Information

Catalog Number: GenBank Accession Number:

83945-5-RR NM 001278182

GeneID (NCBI): Size: CloneNo.: 100ul , Concentration: 1000 $\mu g/ml$ by 8320 241036D9

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source 095936 WB 1:5000-1:50000 Rabbit IF/ICC 1:125-1:500 Full Name:

Isotype: eomesodermin homolog (Xenopus

IgG laevis)

Immunogen Catalog Number: Calculated MW:

AG27634 73 aa

> Observed MW: 67 kDa

Applications

Tested Applications: Positive Controls: WB, IF/ICC, ELISA WB: NK-92 cells, Species Specificity: IF/ICC: SH-SY5Y cells, human, mouse

Background Information

EOMES/TRB2 (also named Eomesodermin/Tbr2) encode T-box transcription factor expressed highly in the intermediate progenitor stage of the developing neuron. During migration of the neurons, radial glia divide and migrate towards the surface of the cerebral cortex, there are three stages of cellular development: radial glia, intermediate progenitors, and postmitotic projection neurons. Radial glia express Pax6, while intermediate progenitor cells express Eomesodermin/Tbr2, and postmitotic projection neurons express Tbr1 (PMID:15634788; PMID:16320258). This process also called neurogenesis, and demonstrated that Eomesodermin/Tbr2 has been $implicated in neurodevelopment. According to the mouse model, several articles showed that {\tt EOMES/TRB2}\ KO$ mice having proliferating cells decreased and having smaller upper cortical layers and a smaller sub ventricular zone in the brain (PMID:18940588). What's more, Eomesodermin/Tbr2 has also been found participating in immune response (PMID:20713880).

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83945-5-RR (EOMES antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using EOMES antibody (83945-5-RR, Clone: 241036D9) at dilution of 1:250 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Biolayer interferometry (BLI) kinetic assays of 83945-5-RR against Human EOMES were performed. The affinity constant is below 1 pM.