

For Research Use Only

CD38 Recombinant antibody

Catalog Number: 84070-5-RR



Basic Information

Catalog Number: 84070-5-RR	GenBank Accession Number: BC007964	Purification Method: Protein A purification
Size: 100ul , Concentration: 1000 µg/ml by Nanodrop;	GeneID (NCBI): 952	CloneNo.: 241246C2
Source: Rabbit	ENSEMBL Gene ID: ENSG00000004468	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	UNIPROT ID: P28907	
	Full Name: CD38 molecule	
	Calculated MW: 300 aa, 34 kDa	
	Observed MW: 40 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : Raji cells, Ramos cells, Daudi cells, Jurkat cells, A549 cells, MOLT-4 cells
Species Specificity: human	

Background Information

CD38, also known as ADP-ribosyl cyclase 1, is a type II transmembrane glycoprotein with a short N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites (PMID: 2319135). The extracellular domain of CD38 has bifunctional enzyme activities that catalyze synthesis of cyclic ADP ribose from nicotinamide adenine dinucleotide (NAD) and hydrolysis of cyclic ADP ribose to adenosine diphosphoribose (PMID: 10636863). CD38 is expressed on a variety of hematopoietic and non-hematopoietic cells and is involved in diverse processes such as generation of calcium-mobilizing metabolites, cell activation, and chemotaxis (PMID: 25938500).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1%BSA**

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84070-5-RR (CD38 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.