For Research Use Only

CD38 Recombinant antibody

Catalog Number:84070-6-RR



Basic Information

Catalog Number: GenBank Accession Number:

84070-6-RR BC007964
Size: GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 952

Nanodrop; ENSEMBL Gene ID:
Source: ENSG00000004468
Rabbit UNIPROT ID:
Isotype: P28907

Full Name: CD38 molecule Calculated MW: 300 aa, 34 kDa Observed MW: 45 kDa Purification Method:

Protein A purfication

CloneNo.: 241246D5

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF/ICC 1:125-1:500

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Species Specificity:

human

IgG

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls

WB: Raji cells, Ramos cells, Jurkat cells, MOLT-4 cells

IHC: human tonsillitis tissue,

IF/ICC: Raji cells,

Background Information

CD38, also known as ADP-ribosyl cyclase 1, is a type II transmembrane glycoprotein with a short N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites (PMID: 2319135). The extracellular domain of CD38 has bifunctional enzyme activities that catalyze synthesis of cyclic ADP ribose from nicotinamide adenine dinucleotide (NAD) and hydrolysis of cyclic ADP ribose to adenosine diphosphoribose (PMID: 10636863). CD38 is expressed on a variety of hematopoietic and non-hematopoietic cells and is involved in diverse processes such as generation of calcium-mobilizing metabolites, cell activation, and chemotaxis (PMID: 25938500).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84070-6-RR (CD38 antibody) at dilution of 1:17000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84070-6-RR (CD38 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84070-6-RR (CD38 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Raji cells using CD38 antibody (84070-6-RR, Clone: 241246D5) at dilution of 1:250 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).

Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 84070-6-RR (CD38 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 84070-6-RR against Human CD38 were performed. The affinity constant is 1.11 nM.