## For Research Use Only

## ATP2A1 Recombinant antibody

Catalog Number:84213-5-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

84213-5-RR BC037354 Protein A purfication GeneID (NCBI): Size: CloneNo.:

100ul , Concentration: 1000  $\mu$ g/ml by 487 241500D12 Nanodrop:

**UNIPROT ID:** Recommended Dilutions: 014983 WB 1:5000-1:50000 Rabbit IHC 1:125-1:500 Full Name:

Isotype: ATPase, Ca++ transporting, cardiac muscle, fast twitch 1

IgG Immunogen Catalog Number: Calculated MW: 1001 aa, 110 kDa AG17944

> Observed MW: 110 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse skeletal muscle tissue, rat skeletal muscle

**Purification Method:** 

IHC: mouse skeletal muscle tissue,

## **Background Information**

ATP2A1 also known as SERCA1, encodes one of the SERCA Ca(2+)-ATPases, which are intracellular pumps located in the sarcoplasmic or endoplasmic reticula of muscle cells. This enzyme catalyzes the hydrolysis of ATP coupled with the translocation of calcium from the cytosol to the sarcoplasmic reticulum lumen and is involved in muscular excitation and contraction. Mutations in ATP2A1 cause some autosomal recessive forms of Brody disease(PMID: 23911890, 10914677).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 84213-5-RR (ATP2A1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 84213-5-RR (ATP2A1 antibody) at dilution of 1:250 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 84213-5-RR against Human ATP2A1 were performed. The affinity constant is 6.90 nM.