For Research Use Only

## Di/Tri-Methyl-Histone H3 (Lys36) Recombinant antibody

Catalog Number:84329-1-RR



**Basic Information** 

Catalog Number: GenBank Accession Number:

84329-1-RR BC066245 GeneID (NCBI): Size:

100ul , Concentration: 1000  $\mu g/ml$  by 8350 Nanodrop: **UNIPROT ID:** Source: P68431 Rabbit Full Name:

Isotype: histone cluster 1, H3a IgG Observed MW: 15 kDa

CloneNo.: 241139G6

**Purification Method:** 

Protein A purfication

Recommended Dilutions: WB 1:2000-1:10000 IF/ICC 1:200-1:800

**Applications** 

**Tested Applications:** WB, IF/ICC, Dot Blot, ELISA

Species Specificity:

human

Positive Controls:

WB: A549 cells, HEK-293 cells, U-87 MG cells, U-251 cells, NIH/3T3 cells, mouse liver tissue, mouse brain

tissue

IF/ICC: HeLa cells,

## **Background Information**

Histones are small, highly basic proteins that consist of a globular domain with unstructured N- and C-terminal tails protruding from the main structure. Histone H3 is one of the five main histones that are responsible for the nucleosome structure of the chromosomal fiber in eukarvotes. Two molecules of each of the four core histones (H2A. H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. In addition to their role in DNA compartmentalization, histones also play crucial roles in various biologic processes, including gene expression and regulation, DNA repair, chromatin condensation, cell cycle progression, chromosome segregation, and apoptosis. The ability of histones to regulate chromatin dynamics primarily originates from various posttranslational modifications carried out by histone-modifying enzymes.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**



Column 1: Unmodified-H3K3f Column 2: H3K36me Column 3: H3K36me3 Column 4: H3K36me3 Column 5: H3K36ac Column 6: Unmodified-H3Y4\* Column 7: H3Y41P Column 7: H3Y41P

Dot blot analysis was used to confirm the specificity of Di/Tri-Methyl-Histone H3 (Lys36) antibody. Acetylated peptides were spotted onto NC and probed with antibody at 1  $\mu$ g/ml.The amount of peptide ( $\mu$ g/mL) spotted is indicated next to each row.



Various lysates were subjected to SDS PAGE followed by western blot with 84329-1-RR (Di/Tri-Methyl-Histone H3 (Lys36) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HIST1H3A antibody (84329-1-RR, Clone: 241139G6) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).