For Research Use Only

DDR1 Recombinant antibody

Catalog Number:84438-6-RR



Basic Information

Catalog Number: GenBank Accession Number: **Purification Method:** 84438-6-RR NM_001297654.2

Protein A purfication GeneID (NCBI): CloneNo.:

100ul , Concentration: 1000 $\mu g/ml$ by 780 241796D10 Nanodrop: **UNIPROT ID:** Recommended Dilutions:

Source Q08345-1 WB 1:500-1:2000 Rabbit IHC 1:50-1:500 Full Name:

Isotype: discoidin domain receptor tyrosine

IgG kinase 1

> Calculated MW: 101kDa Observed MW: 120 kDa

Applications

Tested Applications: Positive Controls:

WB, IHC, ELISA WB: K-562 cells, HeLa cells, HepG2 cells

Species Specificity: IHC: mouse brain tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

DDR1, or Discoidin Domain Receptor 1, is a member of the receptor tyrosine kinase (RTK) family, which plays a significant role in various cellular processes including cell proliferation, adhesion, migration, and extracellular matrix (ECM) remodeling. DDR1 is known for its unique discoidin domain that allows it to bind specifically to certain collagens, initiating downstream signaling pathways that can lead to cell transformation and tumor progression. Abnormal activation of DDR1 is closely associated with the development of various solid tumors, and it has been shown that DDR1 can prevent immune cells from infiltrating triple-negative breast cancer (TNBC) and eliminate tumor cells.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84438-6-RR (DDR1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 84438-6-RR (DDR1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).