## For Research Use Only

## UHRF1 Recombinant antibody

Catalog Number:84489-4-RR Featured Product




Basic Information	Catalog Number: 84489-4-RR	GenBank Accession Number: BC113875	Purification Method: Protein A purfication	
	Size:GenelD (NCBI):100ul, Concentration: 1000 ug/ml by29128Nanodrop;UNIPROT ID:Source:Q96T88RabbitFull Name:Isotype:ubiquitin-like with PHD and ringIgGfinger domains 1Immunogen Catalog Number:Calculated MW:AG17498806 aa, 91 kDaObserved MW:100 kDa		CloneNo.: 241872B12	
			Recommended Dilutions: WB 1:5000-1:50000	
		IF/ICC 1:200-1:800		
		Calculated MW:		
Applications	·		e Controls:	
	WB, IF/ICC, FC (Intra), ELISA Species Specificity: human		WB : MCF-7 cells, HEK-293T cells, HuH-7 cells, HeLa cells, Jurkat cells, HepG2 cells, HCT-116 cells, U2OS cells	
		IF/ICC	HepG2 cells,	
Background Information	UHRF1, also named as ICBP90, NP95 and RNF106, is a putative E3 ubiquitin-protein ligase. It may participate in methylation-dependent transcriptional regulation. UHRF1 binds to inverted 5'-CCAAT-3' box 2 in the TOP2A promoter, and activates TOP2A expression. It is important for G1/S transition and may be involved in DNA repair and chromosomal stability. UHRF1's ability to repress its direct target gene expression is dependent on PHD(UHRF1) binding to unmodified H3R2. In addition, UHRF1 is a novel metabolic guardian restricting AMPK activity(PMID: 34907338).			
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.			
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage		


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84489-4-RR (UHRF1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of UHRF1 antibody (84489-4-RR; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-UHRF1 transfected HEK-293T cells.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using UHRF1 antibody (84489-4-RR, Clone: 241872B12) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10^6 U2OS cells were intracellularly stained with 0.25 ug UHRF1 Recombinant antibody (84489-4-RR, Clone:241872B12) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.