For Research Use Only

VE-cadherin/CD144 Recombinant antibody

Catalog Number:84529-4-RR



Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:5000-1:50000

CloneNo.:

241953A12

Basic Information

Catalog Number: GenBank Accession Number:

84529-4-RR NM 009868.4 GeneID (NCBI):

100ul , Concentration: 1000 $\mu g/ml$ by 12562

Nanodrop: **UNIPROT ID:**

Source: P55284 Rabbit Full Name: Isotype: cadherin 5 IgG Calculated MW:

88kDa Observed MW:

125-130 kDa, 90-100 kDa

Applications

Tested Applications:

WB, ELISA

Species Specificity: mouse

Positive Controls:

WB: bEnd.3 cells,

Background Information

Cadherins are a family of transmembrane glycoproteins that mediate calcium-dependent cell-cell adhesion and play an important role in the maintenance of normal tissue architecture. Vascular endothelial cadherin (VEcadherin), also known as Cadherin-5 (CDH5) or CD144, is a member of the type II classical cadherin family of cell adhesion proteins (PMID: 21269602). VE-cadherin is expressed specifically in endothelial cells and mediates homophilic adhesion in the vascular endothelium (PMID: 1522121; 8555485; 21269602). VE-cadherin plays a role in the organization of lateral endothelial junctions and in the control of permeability properties of vascular endothelium (PMID: 1522121). VE-cadherin has also been shown to be required for angiogenesis (PMID: 16473763; 18162609). The calculated molecular weight of VE-cadherin is 88 kDa and the apparent molecular weight of 120-140 kDa is higher due to post-translational glycosylation and phosphorylation (PMID: 10460833; 29894844). Fulllength VE-cadherin can be proteolytically cleaved to generate a fragment of 90-100 kDa (PMID: 9786462; 22064597).

Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 84529-4-RR (VE-cadherin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Biolayer interferometry (BLL) kinetic assays of 84529-4-RR against Mouse VE-cadherin/CD144 were performed. The affinity constant is 51.6 nM.