For Research Use Only

CD268/BAFF-R Recombinant antibody, PBS Only (Capture)

Catalog Number:84585-4-PBS



Purification Method:

Protein A purification

CloneNo.:

242003C5

Basic Information

Catalog Number: GenBank Accession Number:

84585-4-PBS BC105123

Size: Genel D (NCBI): 100ug, Concentration: 1 mg/ml by 115650

100ug , Concentration: 1 mg/ml by115650Nanodrop;UNIPROT ID:Source:Q96RJ3RabbitFull Name:

Isotype: tumor necrosis factor receptor IgG superfamily, member 13C

> Calculated MW: 184 aa, 19 kDa

Applications

Tested Applications:

Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity:

human

Product Information

84585-4-PBS targets CD268/BAFF-R as part of a matched antibody pair:

MP01419-3: 84585-4-PBS capture and 84585-3-PBS detection (validated in Sandwich ELISA)

Unconjugated rabbit recombinant monoclonal antibody in PBS only (BSA and azide free) storage buffer at a concentration of 1 mg/mL, ready for conjugation. Created using Proteintech's proprietary in-house recombinant technology. Recombinant production enables unrivalled batch-to-batch consistency, easy scale-up, and future security of supply.

This conjugation ready format makes antibodies ideal for use in many applications including: ELISAs, multiplex assays requiring matched pairs, mass cytometry, and multiplex imaging applications. Antibody use should be optimized by the end user for each application and assay.


Background Information

B-cell-activating factor receptor (BAFF-R, also known as CD268 and TNFRSF13C) is a member of the tumor necrosis factor receptor (TNFR) superfamily. It is expressed in most malignant B cells, including those in primary central nervous system lymphoma (PCNSL), and is involved in the survival and proliferation of these cells (PMID: 32528875). The interaction of BAFF with BAFF-R promotes NF-κB activation, which plays a crucial role in B-cell maturation, survival, and costimulation of T-cell activation and proliferation (PMID: 30349534). Additionally, BAFF-R is associated with chronic kidney disease (CKD) by activating NF-κB signaling in renal tubular epithelial cells (PMID: 38791453).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data



Sandwich ELISA standard curve of MP01419-3, Human TNFRSF13C Recombinant Matched Antibody Pair - PBS only. 84585-4-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Eg1922. 84585-3-PBS was HRP conjugated as the detection antibody. Range: 0.156-10 ng/mL



Serum of six individual healthy human donors was measured. The TNFRSF13C concentration of detected samples was determined to be 0.64 ng/mL with a range of 0.27-0.85 ng/mL



Biolayer interferometry (BLL) kinetic assay of 84585-4-PBS against Human TNFRSF13C was performed. The affinity constant is 3.96 nM.