

For Research Use Only

# CD82 Recombinant antibody, PBS Only

Catalog Number: 84617-7-PBS



## Basic Information

**Catalog Number:**

84617-7-PBS

**Size:**

100ug, Concentration: 1 mg/ml by Nanodrop;

**Source:**

Rabbit

**Isotype:**

IgG

**GenBank Accession Number:**

NM\_002231.4

**GeneID (NCBI):**

3732

**UNIPROT ID:**

P27701-1

**Full Name:**

CD82 molecule

**Calculated MW:**

30 kDa

**Observed MW:**

35-45 kDa

**Purification Method:**

Protein A purification

**CloneNo.:**

241765G7

## Applications

**Tested Applications:**

WB, IHC, Indirect ELISA

**Species Specificity:**

human

## Background Information

CD82 is a membrane glycoprotein and belongs to the tetraspanin superfamily, many of which are implicated in the regulation of cell motility, morphology, fusion, signaling, fertilization, and differentiation. CD82 was originally identified as a suppressor of metastasis located on human chromosome 11p11.2 in prostate carcinoma. The majority of evidence indicates that CD82 expression is downregulated or abolished in a variety of malignant tumors. CD82 is present at high levels in human monocyte and macrophage lineages and in various epithelial cells in the prostate, lung, pancreas and many other tissues. In epithelial cells, CD82 is implicated in diverse biological processes such as cell adhesion, migration, apoptosis and morphogenesis.

## Storage

**Storage:**

Store at -80°C.

**Storage Buffer:**

PBS Only


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 84617-7-RR (CD82 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84617-7-PBS in a different storage buffer formulation.




Raji cells were subjected to SDS PAGE followed by western blot with 84617-7-RR (CD82 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84617-7-PBS in a different storage buffer formulation.




U-87 MG cells were subjected to SDS PAGE followed by western blot with 84617-7-RR (CD82 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 84617-7-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 84617-7-RR (CD82 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84617-7-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 84617-7-RR (CD82 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84617-7-PBS in a different storage buffer formulation.



Bi-layer interferometry (BLI) kinetic assays of 84617-7-RR against Human CD82 were performed. The affinity constant is below 1 pM.