

For Research Use Only

# P62,SQSTM1 Recombinant antibody

Catalog Number:84826-1-RR



## Basic Information

<b>Catalog Number:</b> 84826-1-RR	<b>GenBank Accession Number:</b> NM_011018	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ul , Concentration: 1000 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 18412	<b>CloneNo.:</b> 241992C4
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q64337	<b>Recommended Dilutions:</b> WB 1:5000-1:50000
<b>Isotype:</b> IgG	<b>Full Name:</b> sequestosome 1	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
<b>Immunogen Catalog Number:</b> AG35064	<b>Calculated MW:</b> 48 kDa	IHC 1:200-1:800
	<b>Observed MW:</b> 62 kDa	IF/ICC 1:250-1:1000

## Applications

**Tested Applications:**  
WB, IHC, IF/ICC, IP, ELISA

**Species Specificity:**  
human, mouse, rat

**Note-IHC: suggested antigen retrieval with  
TE buffer pH 9.0; (\*) Alternatively, antigen  
retrieval may be performed with citrate  
buffer pH 6.0**

**Positive Controls:**

**WB :** NIH/3T3 cells, HeLa cells, rat brain tissue, HepG2  
cells, Jurkat cells, Raji cells, HEK-293T cells, MCF-7  
cells

**IP :** NIH/3T3 cells,

**IHC :** mouse brain tissue,

**IF/ICC :** NIH/3T3 cells,

## Background Information

Sequestosome 1 (SQSTM1/p62) is a multifunctional adaptor protein implicated in selective autophagy, cell signaling pathways, and tumorigenesis. p62 has been implicated in shuttling ubiquitinated and aggregated proteins for autophagic degradation. p62 is degraded during the autophagic process, which makes its intracellular level a marker for autophagy progression. p62 is at the cross-roads of several signaling pathways including Ras/ Raf/ MAPK and NFκB and plays important role in cancer.

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

**\*\*\* 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech  
Group brand and is not available to purchase from any  
other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62,Sqstm1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




rat brain tissue were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62,SQSTM1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 84826-1-RR (P62,SQSTM1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




IP result of anti-P62,SQSTM1 (IP:84826-1-RR, 4ug; Detection:84826-1-RR 1:4000) with NIH/3T3 cells lysate 1920 ug.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 84826-1-RR (P62,SQSTM1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 84826-1-RR (P62,SQSTM1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using P62,SQSTM1 antibody (84826-1-RR, Clone: 241992C4) at dilution of 1:500 and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Bi-layer interferometry (BLI) kinetic assays of 84826-1-RR against Mouse P62,SQSTM1 were performed. The affinity constant is 0.203 nM.