For Research Use Only

APP Recombinant antibody

Catalog Number:84829-6-RR



Purification Method:

Protein A purification

Basic Information

Catalog Number: GenBank Accession Number:

84829-6-RR NM 000484.3

GeneID (NCBI): CloneNo.: 100ul , Concentration: 1000 $\mu g/ml$ by 351 250546B3

Nanodrop: **UNIPROT ID:** Recommended Dilutions: Source P05067-1 WB: 1:5000-1:10000 Rabbit IF/ICC: 1:200-1:800 Full Name:

Isotype: amyloid beta (A4) precursor protein

IgG Calculated MW:

Immunogen Catalog Number: 87 kDa

EG3070 Observed MW:

100 kDa

Applications

Positive Controls: **Tested Applications:**

WB, IF/ICC, ELISA WB: fetal human brain tissue,

Species Specificity: IF/ICC: U-251 cells, human

Background Information

 $A\beta\ derives\ from\ APP\ via\ proteolytic\ cleavage\ by\ proteases\ called\ a\text{-}, \beta\text{-}\ and\ \gamma\text{-}secretase.\ The\ a\text{-}secretase\ cleavage}$ precludes the formation of A β , while the β - and γ -cleavages generate APP components with amyloidogenic features. Amyloid beta A4 precursor protein(APP), encoded by APP gene which locate on human chromosome 21q, is a cell surface receptor and performs physiological functions on the surface of neurons relevant to neurite growth, neuronal adhesion and axonogenesis. APP expressed in all fetal tissues and is pronounced in brain, kidney, heart and spleen, but weak in liver. Defects in APP are the cause of Alzheimer disease type 1 (AD1). Amyloid β (A β) precursor protein (APP) is a 100-140 kDa transmembrane glycoprotein that exists as several isoforms. APP can be cleaved into several chains, this antibody could recognize N-terminal fragment of APP (N-APP).

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com

Selected Validation Data



fetal human brain tissue were subjected to SDS PAGE followed by western blot with 84829-6-RR (APP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed U-251 cells using APP antibody (84829-6-RR, Clone: 250546B3) at dilution of 1:400 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).