

For Research Use Only

# CNDP2 Recombinant antibody

Catalog Number: 84924-7-RR



## Basic Information

<b>Catalog Number:</b> 84924-7-RR	<b>GenBank Accession Number:</b> BC001375	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 100ul , Concentration: 1000 µg/ml by Nanodrop;	<b>GeneID (NCBI):</b> 55748	<b>CloneNo.:</b> 242373E4
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> Q96KP4	<b>Recommended Dilutions:</b> WB 1:2000-1:10000 IHC 1:1000-1:4000
<b>Isotype:</b> IgG	<b>Full Name:</b> CNDP dipeptidase 2 (metallopeptidase M20 family)	
<b>Immunogen Catalog Number:</b> AG6684	<b>Calculated MW:</b> 53 kDa	
	<b>Observed MW:</b> 44-53 kDa	

## Applications

<b>Tested Applications:</b> WB, IHC, ELISA	<b>Positive Controls:</b>
<b>Species Specificity:</b> human, mouse, rat	<b>WB :</b> LNCaP cells, U2OS cells, mouse kidney tissue, rat kidney tissue
<b>Note-IHC:</b> suggested antigen retrieval with <b>TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	<b>IHC :</b> rat stomach tissue,

## Background Information

Carnosine dipeptidase II (CNDP2) catalyzes the peptide bond hydrolysis in dipeptides, displaying a non-redundant activity toward threonyl dipeptides. It has high dipeptidase activity toward cysteinyl glycine, an intermediate metabolite in glutathione metabolism (PubMed:19346245, PubMed:12473676). Metabolizes N-lactoyl-amino acids, both through hydrolysis to form lactic acid and amino acids, as well as through their formation by reverse proteolysis (PubMed:25964343). CNDP2 has 2 isoforms with a molecular weight of 53 and 44 kDa. The isoform 1 is ubiquitously expressed with higher levels in the kidney and liver (at protein level) and expressed in peripheral blood leukocytes (PubMed:12473676). It is also expressed in gastric mucosa and down-regulated in gastric cancer mucosal tissues (at the protein level) (PubMed:24395568). Isoform 2 is broadly expressed in fetal tissues and expressed in the adult liver and placenta. (PMID: 17121880).

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84924-7-RR (CNDP2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded rat stomach tissue slide using 84924-7-RR (CNDP2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat stomach tissue slide using 84924-7-RR (CNDP2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).