For Research Use Only

CEACAM6 Recombinant antibody

Catalog Number:84982-2-RR



Purification Method:

Basic Information

Catalog Number: GenBank Accession Number:

84982-2-RR BC005008 Protein A purfication GeneID (NCBI): CloneNo.:

100ul , Concentration: 1000 $\mu g/ml$ by 4680 241997D5

Nanodrop; **UNIPROT ID:** Recommended Dilutions: Source: P40199 WB 1:2000-1:10000

Rabbit Full Name:

Isotype: carcinoembryonic antigen-related cell adhesion molecule 6 (non-IgG

specific cross reacting antigen)

Calculated MW: 37 kDa Observed MW: 45-100 kDa

Applications

Tested Applications:

WB, ELISA WB: A549 cells, HT-29 cells

Species Specificity:

human

Background Information

 ${\sf CEACAM6}, also known as \, {\sf CD66c}, is \, a \, gly cosylphosphatidy linositol \, ({\sf GPI})-linked \, cell \, surface \, protein \, that \, belongs \, to \, {\sf CPI}-linked \, cell \, surface \, protein \, that \, belongs \, to \, {\sf CPI}-linked \, cell \, surface \, protein \, that \, belongs \, to \, {\sf CPI}-linked \, cell \, surface \, protein \, that \, belongs \, to \, {\sf CPI}-linked \, cell \, surface \, protein \, that \, belongs \, to \, {\sf CPI}-linked \, cell \, surface \, protein \, that \, belongs \, to \, {\sf CPI}-linked \, cell \, surface \, protein \, that \, belongs \, to \, {\sf CPI}-linked \, cell \, surface \, protein \, that \, {\sf CPI}-linked \, ce$ the CEACAM family of the immunoglobulin superfamily (PMID: 23903773). It is normally expressed on the surface of myeloid and epithelial surfaces and upregulated preferentially at the apical/luminal membranes of many tumors (PMID: 27595061; 35082925). CEACAM6 plays an important role in cell adhesion, proliferation, tumorigenesis, metastasis, and immunity (PMID: 31797958).

Positive Controls:

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 84982-2-RR (CEACAM6 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.