

For Research Use Only

Phospho-PAK1 (Ser144)/PAK2 (Ser141) Recombinant antibody, PBS Only

Catalog Number: 85044-1-PBS



Basic Information

Catalog Number: 85044-1-PBS	GenBank Accession Number: BC109299	Purification Method: Protein A purification
Size: 100ug, Concentration: 1 mg/ml by Nanodrop;	GeneID (NCBI): 5058	CloneNo.: 242123D12
Source: Rabbit	UNIPROT ID: Q13153	
Isotype: IgG	Full Name: p21 protein (Cdc42/Rac)-activated kinase 1	
	Calculated MW: 553 aa, 62 kDa	
	Observed MW: 61 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human, mouse, rat

Background Information

The human PAK family is divided into the group I (PAK1 to PAK3) and group II (PAK4 to PAK6). Group I PAK share some domains that are not present in the group II members. In particular, the autoinhibitory domain (AID) is important for regulation of the kinase activity of the group I family members. Autophosphorylation at PAK1 Ser144, or at the equivalent sites for the other PAK, stabilizes the open conformation and sustains high kinase activity. Mutation of tyrosines 131 or 429 is associated with reduced dimerization and enhanced kinase activity. (PMID: 31748572)

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS Only



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated RAW 264.7 cells and LPS treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 85044-1-RR (Phospho-PAK1 (Ser144)/PAK2 (Ser141) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 85044-1-PBS in a different storage buffer formulation.



Non-treated C6 cells and pervanadate treated C6 cells were subjected to SDS PAGE followed by western blot with 85044-1-RR (Phospho-PAK1 (Ser144)/PAK2 (Ser141) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 85044-1-PBS in a different storage buffer formulation.